

READ SAFETY DIRECTIONS



Amgrow

RotGuard

Systemic Fungicide

ACTIVE CONSTITUENT:
200 g/L PHOSPHOROUS ACID
PRESENT AS MONO-DI POTASSIUM PHOSPHITE
For Citrus, Avocados, Grapes & Ornamentals
Controls Root rot, Collar rot, Downy mildew
Contents 250mL

Downy Mildew

Collar Rot

Root Rot

Amgrow

HOW TO PREPARE: Mix the required rate with water.
CAUTION: DO NOT allow chemical containers or spray to get into drains, sewers, streams or ponds.
STORAGE AND DISPOSAL: Store in the closed, original container in a cool, dry place out of reach of children. DO NOT store in direct sunlight. Dispose of empty container by wrapping in paper, placing in plastic bag and putting in garbage.
SAFETY DIRECTIONS: May irritate the eyes and skin. Avoid contact with eyes and skin. Wash hands after use.
FIRST AID: If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131126.

Amgrow P/L
ABN 61 100 894 786
82 Christensen Road,
Stapledon Qld 4207
Customer Service Number
1800 063 619
www.amgrow.com.au

APVMA
Approval Number:
6675395429

Batch No:

Date of Manufacture:

9310943810972

HOW TO USE: SHAKE WELL BEFORE USE.
DO NOT spray when shade temperatures exceed 35°C.
DO NOT spray when soil is dry and plants are suffering from moisture stress.
Applying as a protective spray will give more effective control. Ensure thorough leaf coverage.

| CROP | DISEASE | RATE | HOW TO APPLY |
|---|---|----------------------------------|---|
| Young or small citrus and recently transplanted trees | Phytophthora Root rot Collar rot | 5-10mL/L | Use higher rate under higher disease risk conditions (e.g. for trees in heavy clay soil). Apply each year as a preventative foliar spray, in late winter prior to flowering and again in autumn when fruit are approaching maturity. Spray until leaves are just wet. Use 5-10 mL/L rate for container grown Mandarin trees. Removal of fruit from affected trees will assist recovery. |
| Mature citrus (except container grown Mandarin trees) | | 100-160 mL in 12L water per tree | |
| Ornamentals | Phytophthora Root rot Collar rot Crown rot | 5 mL/L | Apply at 4-6 weekly intervals as a foliar spray when conditions favour disease. Certain plants may be sensitive to phosphorous acid sprays. Test on a small sample first if in doubt. Do not apply when plants are dormant or stressed. |
| Avocado | Phytophthora Root rot | 2.5-3mL/L | Apply as a foliar spray to run-off. Use 7.5-10 L per adult tree. Apply every 3 weeks as a curative spray, or every 5-6 weeks as a preventative spray. |
| Grapes | Downy Mildew | 6mL/L | Apply as a foliar spray to the point of run-off. Apply at first sign of disease. |

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED

Amgrow

HOW TO PREPARE: Mix the required rate with water.
CAUTION: DO NOT allow chemical containers or spray to get into drains, sewers, streams or ponds.
STORAGE AND DISPOSAL: Store in the closed, original container in a cool, dry place out of reach of children. DO NOT store in direct sunlight. Dispose of empty container by wrapping in paper, placing in plastic bag and putting in garbage.
SAFETY DIRECTIONS: May irritate the eyes and skin. Avoid contact with eyes and skin. Wash hands after use.
FIRST AID: If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131126.

Amgrow P/L
ABN 61 100 894 786
82 Christensen Road,
Stapledon Qld 4207
Customer Service Number
1800 063 619
www.amgrow.com.au

APVMA
Approval Number:
6675395429

Batch No:

Date of Manufacture:

9310943810972