

HOW TO USE: (the measuring spoon provided holds 3gms)

CROP	DISEASE	RATE per 9L of Water	WHP	HOW TO APPLY
ORNAMENTALS				
Ferns	Botrytis leaf spot	9g	-	Apply when disease symptoms first appear and then repeat at 7-10 day intervals.
Flowers, Shrubs	Septoria leaf spot	18g		Apply when disease symptoms first appear and then repeat at 7-14 day intervals
Camellions	Rust, Alternaria leaf spot			Apply when disease symptoms first appear and then repeat at 7-10 day intervals.
Gladioli	Grey mould (Botrytis)			
Chrysanthemums	Septoria leaf spot, Rust			
Roses	Black Spot			
Turf	Fusarium, Rhizoctonia, Helminthosporium	18g in 9 litres of water per 9 sq metres		Apply 9 litres spray per 9 square metres. Repeat each 7 days during susceptible period. Do not feed treated grass clippings to poultry or animals.
FRUIT				
Apples, Pears	Apple Scab (Black spot), Bitter rot, Fly speck, Pear Scab, Ripe fruit spot, Sooty blotch, Target spot	18g	14 days	Spray at 7-14 day intervals following green tip copper spray. Use shorter spray interval during prolonged wet conditions.
Stone fruit	Brown rot, Freckle, Rust Shot hole			Spray at early bloom, mid to full bloom, petal and shuck fall then at intervals of 14 days.
Mangoes	Anthrachnose	18g		Spray at weekly intervals during flowering then monthly until near harvest.
VEGETABLES				
Beans	Rust, Anthracnose	18g	7 days	Spray at first sign of disease. Repeat at 7-10 day intervals.
Capisicum	Target spot			Spray at first sign of disease. Repeat at 7-10 day intervals.
Cabbage, Cauliflower, Broccoli, Brussel sprouts	Alternaria spot, Anthracnose Downy Mildew, Ring spot			Spray at first sign of disease. Repeat at 7-10 day intervals.
Courbits Pumpkins, Melons, Cucumber, Cantaloupe, Zucchini, Squash	Anthrachnose, Downy Mildew, Gummy stem blight, Septoria spot			Spray at first sign of disease. Repeat at 7-10 day intervals while the risk of disease exists.
Lettuce	Anthrachnose, Downy Mildew, Septoria leaf spot		14 days	Spray at first sign of disease. Repeat at 7-10 day intervals. Use shorter intervals during weather conditions favourable to disease development.
Onions & Garlic	Downy mildew, Purple blotch		7 days	
Peas	Downy Mildew			
Potatoes	Early blight (Target spot) Late blight (Irish blight)			
Spinach, Silver Beet, Beetroot	Cercospora leaf spot, Downy mildew			
Tomatoes	Anthrachnose, Early blight (Target spot), grey leaf spot, Late blight (Irish blight), Leaf mould, Phoma rot			Spray at first sign of disease. Repeat at 7-10 day intervals. Do not apply to tomato seedlings before transplanting.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED.
DO NOT PICK APPLES, MANGOES, STONE FRUIT AND LETTUCE FOR 14 DAYS AFTER SPRAYING.
DO NOT PICK VEGETABLES (EXCEPT LETTUCE), MELONS, CANTALOUPE AND TOMATOES FOR 7 DAYS AFTER SPRAYING.

CAUTION
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS



DISEASE CONTROL

MANCOZEB
FUNGICIDE

ACTIVE CONSTITUENT:
750g/KG MANCOZEB

For the control of
certain diseases of Apples,
Mangoes, Vegetables,
Flowers & Turf.



Net Contents 100g

HOW TO PREPARE: Product can be applied through most garden sprayers. Add about half the required volume of water to the sprayer and then add the required amount of product.

The Chemspray measure supplied contains 3 grams.

Mix or agitate and then add the remaining water. Regular agitation during spraying is necessary to avoid product settling out. Do not store mixed product. Only mix what you can use immediately. Thoroughly cover plants with spray ensuring all leaves are wetted. Avoid excessive run-off.

CAUTION: DO NOT feed grass clippings from treated areas to poultry or animals. Dangerous to fish. DO NOT allow chemical containers or spray to get into drains, sewers, streams or ponds.

STORAGE & DISPOSAL: Store in the closed, original container in a cool, dry place out of the reach of children. Do not store in direct sunlight. Dispose of empty container by wrapping in paper, placing in plastic bag and putting in garbage.

SAFETY DIRECTIONS: May irritate the eyes, nose and throat and skin. Avoid contact with eyes and skin. Do not inhale dust or spray mist. Wash hands after use.

FIRST AID: If poisoning occurs, contact a Doctor or Poisons Information Centre. Phone Australia 131126. Avoid giving alcohol.

APVMA APPROVAL NO. 30578/100/0305

Batch No: DOM:



Amgrow Pty Ltd
ABN 81 100 684 786

82 Christensen Road,
Stapylton QLD 4207

Customer Service Number:
1800 063 619

www.amgrow.com.au

MANCOZEB
FUNGICIDE

DISEASE CONTROL

