

Product Name: AMGROW BIN-DIE SELECTIVE LAWN WEEDER READY TO USE  
APVMA Approval No: 59176/129780



|                         |  |
|-------------------------|--|
| Label Name:             | AMGROW BIN-DIE SELECTIVE LAWN WEEDER READY TO USE  |
| Signal Headings:        | POISON<br>KEEP OUT OF REACH OF CHILDREN<br>READ SAFETY DIRECTIONS  |
| Constituent Statements: | ACTIVE CONSTITUENTS: 12 g/L BROMOXYNIL present as N-OCTANOYL ESTER.<br>12 g/L MCPA present as the 2-ETHYLHEXYL ESTER<br>SOLVENT: 20 g/L AROMATIC HYDROCARBONS  |
| Statement of Claims:    | For the control of Bindii & other weeds in Buffalo & other lawns as indicated in the How to Use table.<br><br>(1 L treats 100 Square Metres)<br>(2 L treats 200 Square Metres)<br>(2.5 L treats 250 Square Metres)<br>(5 L treats 500 Square Metres) |
| Net Contents:           | 1L<br>2.5L<br>2L<br>5L   |
| Restrains:              | DO NOT apply on ST varieties of Buffalo grass.   |
| How to Use:             | This section contains file attachment.   |
| Withholding Periods:    |  |
| How to Prepare:         | HOW TO PREPARE: <ul style="list-style-type: none"><li>• Measure lawn area before application to ensure correct dose rate is applied.</li><li>• Shake contents well before using.</li></ul>   |

|  |   |
|--|---|
|  | <ul style="list-style-type: none"> <li>• Attach Hose-on bottle to hose.</li> <li>• Remove plastic plug from the top of the sprayer.</li> <li>• Turn on the tap (the sprayer will immediately start to spraying product). The tap water will mix the product automatically in the correct ratio.</li> <li>• Walk and spray at a steady pace whilst sweeping the spray in a back and forth arc. Overlap treated areas slightly. Spray weeds until foliage is quite wet. Avoid heavy spraying of surrounding lawn grasses.</li> <li>• Stop spraying by removing the bottle from the hose, or by turning off the tap</li> <li>• Re-insert stopper plug on completion.</li> </ul> <p>Weeds will die slowly and effects may take up to 3 weeks to appear.</p> |
|--|---|

|              |  |
|--------------|--|
| Precautions: | CAUTION: Dangerous to fish. DO NOT allow chemical containers or spray to get into drains, sewers, streams or ponds. DO NOT feed treated grass clippings to livestock. DO NOT allow spray to contact or drift onto plants you do not want killed. |
|--------------|--|

|              |  |
|--------------|--|
| Protections: |  |
|--------------|--|

|                       |   |
|-----------------------|---|
| Storage and Disposal: | STORAGE AND DISPOSAL: Store in the closed, original container in a cool, dry place out of the reach of children. Do not store in direct sunlight. Dispose of empty container by wrapping in paper, placing in plastic bag and putting in garbage. |
|-----------------------|---|

|                    |  |
|--------------------|--|
| Safety Directions: | SAFETY DIRECTIONS: Product is poisonous if inhaled or swallowed. Will irritate the eyes and skin. Avoid contact with eyes and skin. Do not inhale spray mist. When preparing spray wear elbow-length PVC gloves and face shield or goggles. If product on skin immediately wash area with soap and water. If product in eyes wash it out immediately with water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each days use wash gloves and face shield or goggles. |
|--------------------|--|

|                         |   |
|-------------------------|---|
| First Aid Instructions: | FIRST AID: If poisoning occurs, contact a Doctor or Poisons Information Centre. Phone Australia 131126. |
|-------------------------|---|

|                     |  |
|---------------------|--|
| First Aid Warnings: |  |
|---------------------|--|

## HOW TO USE

| Situation   | Weeds Controlled   | Mixing Rate                     | How To Apply  |
|---|--|---------------------------------|---|
| Established Couch, Bent, Kikuyu, Paspalum, Buffalo (except ST varieties), Fescue, Zoysia & Rye Lawns and Turf | Bindii (Jo-Jo), Catsear, Clover (not Tas.), Creeping Oxalis, Cudweed, Dandelion, Plantain, Thistles. | Contents (2L) treats 200 Sq. m. | Apply evenly over the area to be treated. Do not mow for 2 days after treatment. Slight browning of grass may occur under hot conditions. More than one application may be required for clover control. |

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED.**