



## Paspalum, Nutgrass & Clover Selective Weed Killer

### HOW TO USE:

- DO NOT apply if rain is expected within 4 hours.
- DO NOT apply in windy conditions.
- DO NOT apply to damaged or stressed turf.
- DO NOT mow for 3 days after spray application.
- DO NOT fertilise for at least 10 days before or after application.
- DO NOT cover spray buffalo or kikuyu lawns - spot spray only.

SITUATION	WEEDS CONTROLLED	RATE COVER SPRAY	RATE SPOT SPRAY	HOW TO APPLY
Couch lawns (except QLD Blue Couch)	Barnyard Grass, Bindii, Capeweed, Catsear, Chickweed, Clovers, Cotula, Cudweed, Dandelion, Duckweed, Fleabane, Lamb's Tongue, Mouse-Eared Chickweed, Mullumbimby Couch, Nutgrass, Paspalum, Scarlet Pimpernel, Summergrass (Crab-grass), Water Couch, White Clover	Mix 170mL per 2L of water and spray over 10m <sup>2</sup>	Mix 90mL per 1L of water	Spray weeds when young and vigorous. When spot spraying thoroughly wet weeds but avoid excessive run-off. Respray in 10 days if necessary.
Bent and Fescue lawns		Mix 85mL per 2L of water and spray over 10m <sup>2</sup>	Mix 45mL in 1L of water	
Buffalo and Kikuyu Lawns		DO NOT COVER SPRAY BUFFALO OR KIKUYU LAWNS	Mix 45mL per 1 L of water	SPOT SPRAY ONLY ON BUFFALO AND KIKUYU LAWNS AS SOME DAMAGE TO LAWNS IS EXPECTED.

NOT TO BE USED FOR ANY PURPOSE OR IN ANY MANNER CONTRARY TO THIS LABEL UNLESS AUTHORISED.

Amgrow Pty Ltd ABN 81 100 684 786  
82 Christensen Road, Stapylton Qld 4207  
Customer Service Number 1800 063 619  
www.amgrow.com.au



**POISON**  
KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS  
BEFORE OPENING OR USING



## Paspalum, Nutgrass & Clover Selective Weed Killer

ACTIVE CONSTITUENTS:  
74g/L DSMA  
24g/L MCPA present as the sodium salt

For control of Paspalum, Nutgrass,  
Mullumbimby Couch, Clovers and  
other weeds in domestic lawns.

Contents 500mL

### HOW TO PREPARE:

Product can be applied with most garden sprayers. For more effective wetting, add wetting agent to final solution. Avoid using fine droplets or applying product in windy conditions as this will increase spray drift. It is important to apply the recommended spray mixture when weeds are young and actively growing. Results may take time to become evident.

### CAUTION:

DO NOT allow spray to contact other plants that you do not want killed. DO NOT allow chemical containers or sprays to get into drains, sewers, streams or ponds. DO NOT feed grass clippings from treated areas to poultry or other animals.

### STORAGE & DISPOSAL:

Store in the closed original container in a cool, dry place out of the reach of children. Do not store in direct sunlight. Dispose of empty container by wrapping in paper, placing in plastic bag and putting in garbage.

### SAFETY DIRECTIONS:

Poisonous if absorbed by skin contact or swallowed. Will irritate the eyes and skin. Avoid contact with the eyes and skin. Do not inhale vapour. Repeated exposure may cause allergic disorder. When opening the container and preparing spray and using the prepared spray wear cotton overalls buttoned to the neck and wrist and a washable hat and elbow-length PVC gloves and face shield or goggles. If product in eyes, wash it out immediately with water. If product on skin, immediately wash area with soap and water. After each day's use wash gloves, face shield or goggles and contaminated clothing.

### FIRST AID:

If poisoning occurs, contact a doctor or Poisons Information Centre (Phone 131 126)

APVMA Approval Number: 67764/56866

Batch No:

Date of Manufacture: