

DIRECTIONS FOR USE RESTRAINTS

DO NOT apply baits to crops
DO NOT exceed 3 m between bait stations for mice, or 9 m for rats.

SITUATION	PEST	RATE	CRITICAL COMMENTS
In and around industrial, commercial, agricultural domestic, and public service buildings, in sewers, animal houses, wharves and around grain terminals.	Rats – <i>Rattus rattus</i> also known as Black rat, Roof, rat, Ship rat or, Fruit rat and <i>Rattus norvegicus</i> also known as the Brown rat or Norway rat (also known as the House mouse). Mice – <i>Mus musculus</i>	3 - 5 STRATAGEM Soft Baits at each bait location 1 - 2 STRATAGEM Soft Baits in each rat hole/burrow 1 - 3 STRATAGEM	Select suitable bait locations, such as in active rat holes/burrows, along rat runs and under rubbish. Try to establish a barrier of bait locations between living and feeding areas. Secure 3 - 5 STRATAGEM Soft Baits at each bait location. STRATAGEM Soft Baits should be placed in areas inaccessible to non-target species, such as in bait trays, inside bait boxes or within enclosed spaces. 1 - 2 STRATAGEM Soft Baits can also be placed within each active rodent hole/burrow. On initial application, an inspection of bait locations after 3 days may be undertaken to determine activity in bait locations and new locations may be selected if necessary. A subsequent inspection may also be undertaken again after 4 days to replace any soft baits which have been eaten. For severe infestations it is recommended that repeat inspections of all bait locations are conducted at 7 day intervals, replacing fresh STRATAGEM Soft Baits only when STRATAGEM Soft Baits have been eaten and until all signs of rat activity have disappeared. Normally only 3 to 4 baiting rounds (pulses) are required for control.
	also known as the House mouse	Soft Baits at each bait location	Mice have very limited home ranges and do not need to drink. They are more difficult to control than rats because their feeding patterns are more erratic. Selection of bait locations is therefore even more important than for rats. Secure 1 - 3 STRATAGEM Soft Baits in areas inaccessible to non-target species, such as in bait trays, inside bait boxes or within enclosed spaces, spaced 2-3 metres apart. Bait locations should be in areas where activity is obvious, particularly where droppings are seen. Inspect bait locations and replace STRATAGEM Soft Baits as for rats until all signs of mouse activity have disappeared.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

GENERAL INFORMATION
STRATAGEM Soft Bait can be used both indoors and outdoors. Rats and mice that are resistant to warfarin are controlled and no prebaiting is required. A lethal dose is readily consumed in a single feed but death may not occur for up to seven days. DO NOT place bait on or near exposed food or food utensils, food preparation areas or food storage areas.

GENERAL INSTRUCTIONS

Place STRATAGEM Soft Baits in areas frequented by the target pests. DO NOT place baits in areas accessible to domestic animals, children, pets and wildlife. DO NOT place baits in areas liable to flooding. Alternative food sources should be removed or reduced as much as possible.

PRECAUTIONS

Place STRATAGEM Soft Baits in areas frequented by the target pests. DO NOT place baits in areas accessible to domestic animals, children, pets and wildlife. DO NOT place baits in areas liable to flooding. Alternative food sources should be removed or reduced as much as possible.
DO NOT re-handle product unless wearing chemical resistant gloves.
DO NOT place baits in the open.
DO NOT allow baits to contaminated foodstuffs or food intended for human or animal consumption.
DO NOT remove bait from sachet. DO NOT use in areas accessible to children.

POISON
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Stratagem® Soft Bait
Rodenticide



ACTIVE CONSTITUENT: 0.05 g/kg FLOCOUMAFEN

READY TO USE BAIT – KILLS IN A SINGLE FEED

Controls rats and mice in and around industrial, commercial, agricultural and domestic buildings.
Controls rats and mice resistant to warfarin.

IMPORTANT: READ THE LABEL BEFORE OPENING OR USING

BASF Australia Ltd ABN 62 008 437 867
Level 12, 28 Freshwater Place, Southbank VICTORIA 3006
www.crop-solutions.basf.com.au

Net Contents:
2.4 kg



PROTECTION OF LIVESTOCK, WILDLIFE, FISH, CRUSTACEANS AND OTHERS
This bait is hazardous to pigs, cats, dogs, poultry, birds and wildlife. DO NOT place baits in locations which are accessible to other animals or birds. DO NOT contaminate ponds, streams, rivers or waterways with the bait or its used containers. The product is restricted to use in and around buildings (within 2 metres) or enclosed spaces, e.g., drains. In enclosed roof spaces and areas where possums and other non-target animals frequent, baits should be secured inside proper bait stations, plastic jars or plastic tubes. The use of this product for the control of protected native rodents requires permission from wildlife authorities.

STORAGE AND DISPOSAL
Store in the closed original container in a safe place. DO NOT store for prolonged periods in direct sunlight. Containers which have held bait should not be used for any other purpose. Pest Control Operators should dispose of empty containers, used bait stations and unused or untaken bait by burying in an approved dump. In a domestic situation, minimise the presence of excess baits. Where unused baits are in good condition, return to original container for reuse. For empty containers, damaged or unwanted baits, wrap securely and dispose of in household refuse bin. Look for dead rats or mice and burn or bury them. DO NOT place dead rats or mice in refuse bins or rubbish tips. Ensure baits are removed when rodent activity has ceased.

SAFETY DIRECTIONS
May irritate eyes and skin. Avoid contact with eyes and skin. When using the product, wear elbow-length chemical resistant gloves. Wash hands after use. After each day's use, wash gloves.

FIRST AID
If poisoning occurs, contact a doctor or Poisons Information Centre. Telephone 13 11 26 Australia-wide. New Zealand 0800 764 766. Vitamin K (Phytomenadione) is antidotal.

SDS
Additional information is listed in the Safety Data Sheet.

ADDITIONAL STATEMENTS (required by WHS REGULATIONS 2011)
HAZARD STATEMENT: May cause damage to organs (Blood) through prolonged or repeated exposure.
PRECAUTIONARY STATEMENT (Prevention): Do not breathe dust.
PRECAUTIONARY STATEMENT (Response): Get medical advice/attention if you feel unwell.

CONDITIONS OF SALE
All conditions and warranties rights and remedies implied by law or arising in contract or tort whether due to the negligence of BASF Australia Ltd ABN 62008437867 or otherwise are hereby expressly excluded so far as the same may legally be done provided however that any rights of the Buyer pursuant to non-excludable conditions or warranties of the Competition and Consumer Act 2010 or any relevant legislation of any State are expressly preserved but the liability of BASF Australia Ltd or any intermediate Seller pursuant thereto shall be limited if so permitted by the said legislation to the replacement of the goods sold or the supply of equivalent goods and all liability for indirect or consequential loss or damage of whatsoever nature is expressly excluded. This product must be used or applied strictly in accordance with the instructions appearing hereon. This product is solely sold for use in Australia and must not be exported without the prior written consent of BASF Australia Ltd.

APVMA Approval No: 90839/130175
Batch No:
Date of Manufacture:
© STRATAGEM is a registered trademark of BASF
© Copyright, 2021
BASF Australia Ltd
ABN 62 008 437 867
Level 12, 28 Freshwater Place
Southbank VICTORIA 3006
W103684 05.2021

FOR SPECIALIST ADVICE IN AN
EMERGENCY ONLY PHONE 1800 803 440
TOLL FREE – ALL HOURS – AUSTRALIA WIDE

